

Intermediate Algebra Formulas Quiz 2 – Use This to Study

Some factoring formulas:

Sum of Cubes $a^3 + b^3 = (a + b)(a^2 - ab + b^2)$

Difference of Cubes $a^3 - b^3 = (a - b)(a^2 + ab + b^2)$

Difference of Squares $a^2 - b^2 = (a + b)(a - b)$

Some work times formulas:

Two laborers:

$$\frac{1}{x} = \frac{1}{a} + \frac{1}{b}$$

Three laborers:

$$\frac{1}{x} = \frac{1}{a} + \frac{1}{b} + \frac{1}{c}$$

When something moves a distance d at rate r for time t , then $d = rt$

When simplifying a radical: If your variable could be negative and the index of the radical is even the result must be placed in absolute value bars

If n is a positive integer greater than 1 and $\sqrt[n]{a}$ is a real number, then $a^{1/n} = \sqrt[n]{a}$

If m and n are positive integers greater than 1 with $\frac{m}{n}$ in simplest form, then $a^{m/n} = \sqrt[n]{a^m} = (\sqrt[n]{a})^m$

As long as $a^{m/n}$ is a nonzero real number, $a^{-m/n} = \frac{1}{a^{m/n}}$

The Product Rule for Radicals: If $\sqrt[n]{a}$ and $\sqrt[n]{b}$ are real numbers, then $\sqrt[n]{a} \cdot \sqrt[n]{b} = \sqrt[n]{ab}$

The Quotient Rule for Radicals: If $\sqrt[n]{a}$ and $\sqrt[n]{b}$ are real numbers, and $\sqrt[n]{b}$ is not zero, then $\frac{\sqrt[n]{a}}{\sqrt[n]{b}} = \sqrt[n]{\frac{a}{b}}$

The distance between two points, (x_1, y_1) and (x_2, y_2) , is $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

The midpoint of the line segment whose endpoints are (x_1, y_1) and (x_2, y_2) is the point with coordinates $\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$

The Pythagorean Theorem: If a and b are the legs of a right triangle, and c is its hypotenuse, then $a^2 + b^2 = c^2$

The imaginary unit, written i , is the number whose square is -1 . That is, $i^2 = -1$ and $i = \sqrt{-1}$

If a is a positive number, then $\sqrt{-a} = i \cdot \sqrt{a}$

The Square Root Property: If X is any algebraic expression, c is a real number, and $X^2 = c$, then $X = \pm\sqrt{c}$

The Quadratic Formula: If $ax^2 + bx + c = 0$ then $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ and its discriminant $b^2 - 4ac$ tells the quantity and type of solution(s).

Given the parabola $y = f(x) = ax^2 + bx + c$, its vertex is the point $\left(\frac{-b}{2a}, c - \frac{b^2}{4a}\right)$