ANSWERS: Practice Quiz 10 Part A Anti-depressants DEH 2300 - July, 2014

1.	Prozac® (fluoxetine) is considered first line treatment for which disease/condition?										
	a.	ADHD			c.		ng cessatio				
	b.	Depression	ANSWEF	₹	d.	Appeti	ite suppress	sion			
2.	Which condition is likely to occur if a patient taking a drug classified as a monamine oxidase inhibitor consumes										
		foods or beverages such as red wine, aged cheese or organ meats?									
	a.	Severe hyperg			c.		ral edema				
	b.	Hypertensive	crisis A	NSWER	d.	Ulcera	tive colitis				
3.	Which medication is used as an anti-depressant and to assist with cessation of smoking?										
	a.	Wellbutrin® (bupropion) ANSWER					® (alprazo				
	b.	Loniten® (minoxidil) d. Ritalin® (methylphenidate)									
4.	Which group of antidepressants is currently considered first line treatment for depression?										
	a.										
	b.	b. SSRIs (selective serotonin re-uptake inhibitors) ANSWER									
	c.	· · · · · · · · · · · · · · · · · · ·									
	d.	Miscellaneous	S								
5.	Tyramine rich foods are contraindicated with which group of antidepressants?										
	a.	TCAs (tricyclics)									
	b.	SSRIs (selective serotonin re-uptake inhibitors)									
	c.	·									
	d.	Miscellaneous	8								
6.	What	What is the likely outcome of the contraindication in question 5?									
	ANSWER: Extreme hypertension which may become life-threatening if untreated										
7.	Which group of antidepressants is considered "first generation"?										
	a.										
	b.	MAOIs (monoamide oxidase inhibitors)									
	c.	SSRIs (selective serotonin re-uptake inhibitors)									
	d.	Miscellaneous		•	ĺ						
8.	Prozac® (fluoxetine) and Zoloft® (sertraline) are classified as:										
	a.	_` ′									
	b.	MAOIs (monoamide oxidase inhibitors)									
	c.	SSRIs (selective serotonin re-uptake inhibitors) ANSWER									
	d.	Miscellaneous	5	•		ŕ					
9.	Which drug is primarily an antidepressant but has unlabeled uses of treatment for ADHD, neuropathic pain,										
	fibromyalgia, stress incontinence, aggressive behavior and alcohol withdrawal?										
	a. Wellbutrin® (bupropion) ANSWER c. Zoloft® (sertraline)										
	b.	Prozac® (fluo		,	d.		nil® (imipra	,			
10.	Why is Catapres® (clonidine) contraindicated with tricyclics? What is the primary indication for Catapres®?										
	ANS		Hypertensive crisis is very likely to occur. The primary indication for Catapres® is treatment of hypertension.								

11. List at least 3 contraindications for SSRIs?

ANSWER: suicide prone, bipolar disorders, liver impairment, renal impairment, diabetes, eating disorders, pregnancy and lactation

12. Write a brief drug profile for Viibryd® (vilazodone hydrochloride).

ANSWER: See the website below.

http://www.rxlist.com/viibryd-drug.htm